arXiv:1609.01005v1 [math.PR] 5 Sep 2016

The third moment for the parabolic Anderson model

Le Chen*

University of Kansas

Tuesday 6th September, 2016

Abstract

In this paper, we study the *parabolic Anderson model* starting from the Dirac delta initial data:

$$\left(\frac{\partial}{\partial t} - \frac{\nu}{2}\frac{\partial^2}{\partial x^2}\right) u(t,x) = \lambda u(t,x)\dot{W}(t,x), \qquad u(0,x) = \delta_0(x), \quad x \in \mathbb{R},$$

where $\dot{W}$ denotes the space-time white noise. By evaluating the threefold contour integral in the third moment formula by Borodin and Corwin [2], we obtain some explicit formulas for $\mathbb{E}[u(t,x)^3]$. One application of these formulas is given to show the exact phase transition for the intermittency front of order three.

Keywords. The stochastic heat equation; parabolic Anderson model; Dirac delta initial condition; space-time white noise; moment formula; intermittency fronts; growth indices.

AMS 2010 subject classification. Primary 60H15; Secondary 35R60.

1 Introduction

In this paper, we derive an explicit formula for the third moment of the following *parabolic Anderson model* (PAM [3]),

$$\begin{cases} \left(\dfrac{\partial}{\partial t} - \dfrac{\nu}{2}\dfrac{\partial^2}{\partial x^2}\right) u(t,x) = \lambda u(t,x)\, \dot{W}(t,x), \quad x \in \mathbb{R},\ t > 0, \\ u(0,\cdot) = \delta_0, \end{cases} \tag{1.1}$$

where $\nu > 0$ is the diffusion parameter and δ_0 is the Dirac delta measure with a unit mass at zero. The noise is assumed to be the space-time white noise, i.e.,

$$\mathbb{E}\left[\dot{W}(t,x)\dot{W}(s,y)\right] = \delta_0(t-s)\delta_0(x-y).$$

*University of Kansas, Department of Mathematics. 405 Snow Hall, 1460 Jayhawk Blvd. Lawrence, Kansas, 66045-7594, USA. *Email*: chenle@ku.edu or chenle02@gmail.com.

The solution to (1.1) is understood in the mild form

$$u(t,x) = G_\nu(t,x) + \lambda \int_0^t \int_{\mathbb{R}^d} G(t-s, x-y) u(s,y) W(\mathrm{d}s, \mathrm{d}y), \tag{1.2}$$

where the stochastic integral in (1.2) is in the sense of Walsh [13] and

$$G_\nu(t,x) = (2\pi\nu t)^{-1/2} \exp\left(-\frac{x^2}{2\nu t}\right). \tag{1.3}$$

Explicit formulas both for the second moment and for the two-point correlation function were obtained in [4, 7] for general initial measure $u(0,\cdot) = \mu(\cdot)$ such that $(|\mu| * G(t,\circ))(x) < \infty$ for all $t > 0$ and $x \in \mathbb{R}$. Here, $\mu = \mu_+ - \mu_-$ is the *Jordan* decomposition and $|\mu| = \mu_+ + \mu_-$. When the initial data is the delta measure with a unit mass at zero, i.e., $\mu = \delta_0$, these formulas reduce to the following simple forms (see Corollary 2.8 of [4]):

$$\begin{aligned}\mathbb{E}\left[u(t,x)u\left(t,y\right)\right] = G_\nu(t,x)G_\nu\left(t,y\right) + \frac{\lambda^2}{4\nu} G_{\nu/2}\left(t, \frac{x+y}{2}\right) \\ \times \exp\left(\frac{\lambda^4 t - 2\lambda^2|x-y|}{4\nu}\right) \operatorname{erfc}\left(\frac{|x-y| - \lambda^2 t}{2\sqrt{\nu t}}\right),\end{aligned} \tag{1.4}$$

and in particular,

$$\mathbb{E}\left[u(t,x)^2\right] = G_{\nu/2}(t,x)\left(\frac{\lambda^2}{\sqrt{4\pi\nu t}} + \frac{\lambda^4}{2\nu} e^{\frac{\lambda^4 t}{4\nu}} \Phi\left(\lambda^2\sqrt{\frac{t}{2\nu}}\right)\right). \tag{1.5}$$

In these formulas, we have used the following special functions

$$\Phi(x) = \int_{-\infty}^x (2\pi)^{-1/2} e^{-y^2/2} \mathrm{d}y, \quad \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-y^2} \mathrm{d}y, \quad \operatorname{erfc}(x) = 1 - \operatorname{erf}(x).$$

Bertini and Cancrini obtained the two-point correction function in an integral form; see Corollary 2.5 in [1]. This integral has been evaluated explicitly in [4], which gives the same form as (1.4).

On the other hand, for the delta initial condition, in a beautiful work by Borodin and Corwin [2], it is showed that for any $x_1 \le \cdots \le x_k$,

$$\mathbb{E}\left[\prod_{j=1}^k u(t,x_j)\right] = \frac{1}{(2\pi\imath)^k} \int \cdots \int \prod_{1\le A<B\le k} \frac{z_A - z_B}{z_A - z_B - \frac{\lambda^2}{\nu}} \prod_{j=1}^k \exp\left(\frac{\nu t}{2} z_j^2 + x_j z_j\right) \mathrm{d}z_j, \tag{1.6}$$

where $\imath = \sqrt{-1}$ and the z_j integration is over $\alpha_j + \imath\mathbb{R}$ with

$$\alpha_1 > \alpha_2 + \frac{\lambda^2}{\nu} > \alpha_3 + \frac{2\lambda^2}{\nu} > \cdots; \tag{1.7}$$

see Appendix A.2 in [2] where they assume $\nu = 1$. Thanks to condition (1.7), this formula (1.6) can be transformed into the following form by introducing another $k(k-1)/2$ integrals, which results in a formula with $k(k+1)/2$ integrals:

$$\begin{aligned}\mathbb{E}\left[\prod_{j=1}^k u(t,x_j)\right] = \frac{1}{(2\pi\imath)^k} \int \cdots \int \prod_{j=1}^k \exp\left(\frac{\nu t}{2} z_j^2 + x_j z_j\right) \mathrm{d}z_j \\ \times \prod_{1\le A<B\le k} (z_A - z_B) \int_0^\infty \exp\left(-s_{AB}\left[z_A - z_B - \frac{\lambda^2}{\nu}\right]\right) \mathrm{d}s_{AB}.\end{aligned} \tag{1.8}$$

In the following, we will first show that when $k=2$, one can recover (1.4) by evaluating the double contour integrals in (1.6). Then we proceed to derive some formulas, more explicit than (1.6), for the third moment. As an application, we establish the third exact phase transition (see below). These results are summarized in the following three Theorems 1.1, 1.2 and 1.7.

Theorem 1.1 (Second moment). *When $k=2$, the double contour integrals on the right-hand side of (1.6) are equal to*

$$G_\nu(t,x_1)G_\nu(t,x_2)+\frac{\lambda^2}{4\nu}G_{\nu/2}\left(t,\frac{x_1+x_2}{2}\right)\exp\left(\frac{t\lambda^4-2\lambda^2(x_2-x_1)}{4\nu}\right)\operatorname{erfc}\left(\frac{(x_2-x_1)-t\lambda^2}{2\sqrt{\nu t}}\right),$$

which recovers (1.4) (with $x_1\le x_2$).

Note that in case of $k=3$, there are six integrals in (1.8). In the next theorem, we will first evaluate the three contour integrals in (1.8) which leads to (1.9). Then we proceed to evaluate two real integrals in (1.9) leading to (1.10). Finally, by applying the mean-value theorem, we evaluate the last real integral to obtain an explicit expression which is handy for applications; see (1.12).

Theorem 1.2 (Third moment). *Suppose that the initial data is the Dirac delta measure $\delta_0(x)$. The following statements are true:*
(1) For all $t>0$ and $x_i\in\mathbb{R}$, $i=1,2,3$,

$$\begin{aligned}
&\mathbb{E}\left[u(t,x_1)u(t,x_2)u(t,x_3)\right]\\
=&(\nu t)^{-9/2}(2\pi)^{-3/2}\int_0^\infty \mathrm{d}s_1\,\exp\left(\frac{\lambda^2 s_1}{\nu}\right)\int_0^\infty \mathrm{d}s_2\,\exp\left(\frac{\lambda^2 s_2}{\nu}\right)\int_0^\infty \mathrm{d}s_3\,\exp\left(\frac{\lambda^2 s_3}{\nu}\right)\\
&\times\left(-s_1+s_2+2s_3+|x_3-x_2|\right)\left(s_1+2s_2+s_3+|x_3-x_1|\right)\left(2s_1+s_2-s_3+|x_2-x_1|\right)\\
&\times\exp\left(-\frac{(x_3+s_2+s_3)^2}{2\nu t}-\frac{(x_2+s_1-s_3)^2}{2\nu t}-\frac{(x_1-s_1-s_2)^2}{2\nu t}\right).
\end{aligned}\tag{1.9}$$

(2) For all $x\in\mathbb{R}$,

$$\begin{aligned}
\mathbb{E}[u(t,x)^3]=&\frac{\lambda^2}{2^{5/2}\pi\,\nu^3t^2}\exp\left(\frac{\lambda^2 t}{4\nu}-\frac{3x^2}{2\nu t}\right)\\
&\times\int_0^\infty\left[\left(3s+\lambda^2t\right)\exp\left(-\frac{3s(s-2\lambda^2t)}{4\nu t}\right)\Phi\left(\frac{\lambda^2t+s}{\sqrt{2\nu t}}\right)\right.\\
&\qquad\left.+\left(3s-\lambda^2t\right)\exp\left(-\frac{s(3s-2\lambda^2t)}{4\nu t}\right)\Phi\left(\frac{\lambda^2t-s}{\sqrt{2\nu t}}\right)\right]\mathrm{d}s\\
&+\frac{1}{(2\pi\nu t)^{3/2}}\exp\left(-\frac{3x^2}{2\nu t}\right)\\
&+\frac{\lambda^2}{2\sqrt{2}\pi\nu^2t}\exp\left(\frac{\lambda^4t}{4\nu}-\frac{3x^2}{2\nu t}\right)\left(1-\Phi\left(-\lambda^2\sqrt{\frac{t}{2\nu}}\right)\right).
\end{aligned}\tag{1.10}$$

(3) For all $t>0$ and $x\in\mathbb{R}$, there are some two constants a,b depending on t, ν and λ in the following range

$$0\le a\le\Phi\left(\lambda^2\sqrt{\frac{t}{2\nu}}\right)\le b\le 1\tag{1.11}$$

such that

$$\mathbb{E}\left[u(t,x)^3\right] = \frac{1}{(2\pi\nu t)^{3/2}} \exp\left(-\frac{3x^2}{2\nu t}\right) \tag{1.12a}$$

$$+ \frac{\lambda^2(b-a)}{2\sqrt{2}\pi\nu^2 t} \exp\left(\frac{\lambda^4 t}{4\nu} - \frac{3x^2}{2\nu t}\right) \tag{1.12b}$$

$$+ \frac{\lambda^2}{2\sqrt{2}\pi\nu^2 t} \exp\left(\frac{\lambda^4 t}{4\nu} - \frac{3x^2}{2\nu t}\right) \Phi\left(\lambda^2\sqrt{\frac{t}{2\nu}}\right) \tag{1.12c}$$

$$+ \frac{b\sqrt{2}\,\lambda^4}{\sqrt{3\pi\nu^5 t}} \exp\left(\frac{\lambda^4 t}{\nu} - \frac{3x^2}{2\nu t}\right) \Phi\left(\lambda^2\sqrt{\frac{3t}{2\nu}}\right). \tag{1.12d}$$

Remark 1.3. It is known that $u(t,x)$ is strictly positive for all $t > 0$ and $x \in \mathbb{R}$ a.s.; see [6]. Hence, $\mathbb{E}\left[|u(t,x)|^3\right] = \mathbb{E}\left[u(t,x)^3\right]$.

In the following, let u_δ and u_1 denote the solutions to (1.1) starting from the delta measure δ_0 and Lebesgue's measure (i.e., $u(0,x) = 1$), respectively. The following corollary is a consequence of (1.9) and the fact that

$$\mathbb{E}\left[u_1(t,x)^k\right] = \int_{\mathbb{R}} \cdots \int_{\mathbb{R}} \mathbb{E}\left[\prod_{i=1}^{k} u_\delta(t,x_i)\right] \mathrm{d}x_1 \ldots \mathrm{d}x_k. \tag{1.13}$$

Corollary 1.4. *Suppose that the initial data is Lebesgue's measure, i.e., $u(0,x) = 1$. Then*

$$\begin{aligned}
\mathbb{E}\left[u_1(t,x)^3\right] =& (\nu t)^{-9/2}(2\pi)^{-3/2} \int_{\mathbb{R}} \mathrm{d}x_1 \int_{\mathbb{R}} \mathrm{d}x_2 \int_{\mathbb{R}} \mathrm{d}x_3 \\
&\times \int_0^\infty \mathrm{d}s_1 \, \exp\left(\frac{\lambda^2 s_1}{\nu}\right) \int_0^\infty \mathrm{d}s_2 \, \exp\left(\frac{\lambda^2 s_2}{\nu}\right) \int_0^\infty \mathrm{d}s_3 \, \exp\left(\frac{\lambda^2 s_3}{\nu}\right) \\
&\times (-s_1 + s_2 + 2s_3 + |x_3 - x_2|) \\
&\times (\quad s_1 + 2s_2 + s_3 + |x_3 - x_1|) \\
&\times (\quad 2s_1 + s_2 - s_3 + |x_2 - x_1|) \\
&\times \exp\left(-\frac{(x_3 + s_2 + s_3)^2}{2\nu t} - \frac{(x_2 + s_1 - s_3)^2}{2\nu t} - \frac{(x_1 - s_1 - s_2)^2}{2\nu t}\right).
\end{aligned} \tag{1.14}$$

Remark 1.5. Bertini and Cancrini studied (1.1) with $\lambda = 1$ and claimed in Theorem 2.6 of [1] that

$$\mathbb{E}\left[u_1(t,x)^k\right] = 2\exp\left(\frac{k(k^2-1)}{4!\,\nu} t\right) \Phi\left(\sqrt{\frac{k(k^2-1)}{12\nu}} t\right). \tag{1.15}$$

X. Chen showed in [8] that (1.15) is correct only for $k = 2$; see also Remark 2.6 in [4]. Note that there are six integrals in (1.14). After integrals over $\mathrm{d}x_2\mathrm{d}x_3$, the expression becomes too complicated and too long to handle. We leave it to interested readers to simplify this formula.

Remark 1.6 (Asymptotics)**.** Note that the leading orders for large time t both in (1.15) with $k = 3$ for u_1 and in (1.12) for u_δ are the same. Actually, X. Chen [8] showed that the asymptotics of the k-th moment of u_1 (note that it is not u_δ) [1] does satisfy, as (1.15), that

$$\lim_{t\to\infty} \frac{1}{t} \log \mathbb{E}\left[u_1(t,x)^k\right] = \frac{k(k^2-1)}{4!\,\nu}. \tag{1.16}$$

As an easy consequence of (1.12), we see that this fact is still true for u_δ, namely,

$$\lim_{t\to\infty} \frac{1}{t} \log \mathbb{E}\left[u_\delta(t,x)^3\right] = \frac{\lambda^4 k(k^2-1)}{4!\,\nu}\bigg|_{k=3} = \frac{\lambda^4}{\nu}, \qquad \text{for all } x \in \mathbb{R}. \tag{1.17}$$

Of course, the formula (1.12) itself contains more information than these asymptotics.

Now we state one application of these moment formulas. It is known that the solution to (1.1) with a general initial condition is intermittent [3, 4, 10], which means informally that the solution in question develops many tall peaks. An interesting phenomenon is that when the initial data has compact support, these tall peaks will propagate in space with certain speed depending on the value of λ; see some simulations in Figure 1. The spatial fronts of these tall peaks are called the *intermittency fronts*. Conus and Khoshnevisan [9] introduced the following *lower and upper growth indices of order p* to characterize these intermittency fronts,

$$\underline{\lambda}(p) := \sup\left\{\alpha > 0 : \limsup_{t\to\infty} \frac{1}{t} \sup_{|x|\ge \alpha t} \log \mathbb{E}\left(|u(t,x)|^p\right) > 0\right\}, \tag{1.18}$$

$$\overline{\lambda}(p) := \inf\left\{\alpha > 0 : \limsup_{t\to\infty} \frac{1}{t} \sup_{|x|\ge \alpha t} \log \mathbb{E}\left(|u(t,x)|^p\right) < 0\right\}. \tag{1.19}$$

We call the case $\underline{\lambda}(p) = \overline{\lambda}(p)$ the *p-th exact phase transition*. Chen and Dalang [4] established the second exact phase transition, namely, if the initial data is a nonnegative measure with compact support, then

$$\lambda(2) := \underline{\lambda}(2) = \overline{\lambda}(2) = \frac{1}{2}\,\lambda^2. \tag{1.20}$$

This improves the result by Conus and Khoshnevisan [9] that $(2\pi)^{-1}\lambda^2 \le \underline{\lambda}(2) \le \overline{\lambda}(2) \le 2^{-1}\lambda^2$. Chen and Kunwoo [5] studied the stochastic heat equation – the equation with λu in (1.1) replaced by $\sigma(u)$ – on $\mathbb{R}^d$ subject to a Gaussian noise that is white in time and colored in space. Both nontrivial lower and upper bounds for the second growth indices were obtained. More recently, Huang, Lê and Nualart studied the PAM on $\mathbb{R}^d$ with a Gaussian noise that may have colors in both space and time; see [11] and [12]. They obtained some nontrivial, sometimes matching, bounds for the lower and upper growth indices of all orders $p \ge 2$. In particular, they showed in the current setting that

$$\lambda(p) = \underline{\lambda}(p) = \overline{\lambda}(p) = \sqrt{\frac{p^2-1}{12}}\,\lambda^2. \tag{1.21}$$

[1] Note that in [8], the initial data is assumed to a nonnegative function that satisfies

$$0 < \inf_{x\in\mathbb{R}} u(0,x) \le \sup_{x\in\mathbb{R}} u(0,x) < \infty.$$

For studying asymptotic properties, this assumption is essentially equivalent to the case that $u(0,x) \equiv 1$.

The following theorem is an easy corollary of our moment formula, which recovers their result (1.21) for $p = 3$.



(a) $\lambda = 3$ (b) $\lambda = 4$ (c) $\lambda = 5$

(d) $\lambda = 6$ (e) $\lambda = 7$ (f) $\lambda = 8$

Figure 1: Some simulations of the solution to (1.1) with various λ ranging from 3 to 8. These simulations are truncated at the level 500 in order to compare the size of these six space-time cones. The initial data is $(2\pi\mu)^{-1/2} \exp(-x^2/(2\mu))$ with $\mu = 10^{-5}$, which is an approximation to the delta initial measure. These simulations are made up to $t = 0.01$.

Theorem 1.7 (The third exact phase transition). *If* $u(0, x) = \delta_0(x)$, *then*

$$\lambda(3) := \overline{\lambda}(3) = \underline{\lambda}(3) = \sqrt{\frac{2}{3}}\,\lambda^2. \tag{1.22}$$

Proof. Because b in (1.11) goes to 1 as t goes to infinity, we see that

$$\lim_{t\to\infty} \frac{1}{t} \sup_{|x|\geq \alpha t} \log \mathbb{E}\left(u(t,x)^p\right) = \frac{\lambda^4}{\nu} - \frac{3\alpha^2}{2\nu}. \tag{1.23}$$

Hence,

$$\lim_{t\to\infty} \frac{1}{t} \sup_{|x|\geq \alpha t} \log \mathbb{E}\left(u(t,x)^p\right) > 0 \qquad \Longleftrightarrow \qquad \alpha < \sqrt{\frac{2}{3}}\,\lambda^2, \tag{1.24}$$

which shows that $\underline{\lambda}(3) = \sqrt{2/3}\,\lambda^2$. Similarly, one can show that $\overline{\lambda}(3) = \sqrt{2/3}\,\lambda^2$. □

In the following, we will prove Theorems 1.1 and 1.2 in Sections 2 and 3, respectively.

2 Second moment: Proof of Theorem 1.1

The proof of Theorem 1.1 severs as a warm-up for the more involved proof of Theorem 1.2.

Proof of Theorem 1.1. Suppose that $x_1 \le x_2$. Fix any $\alpha_1 > \alpha_2 + \frac{\lambda^2}{\nu}$. Then, from (1.8),

$$\begin{aligned}\mathbb{E}\left[u(t,x_1)u(t,x_2)\right] =& \frac{1}{(2\pi\imath)^2}\iint\left(\int_0^\infty \exp\left(-\left[(z_1-z_2)-\frac{\lambda^2}{\nu}\right]s\right)\mathrm{d}s\right)\\ &\times(z_1-z_2)\exp\left(\frac{\nu t}{2}z_1^2+x_1z_1\right)\exp\left(\frac{\nu t}{2}z_2^2+x_2z_2\right)\mathrm{d}z_1\mathrm{d}z_2\\ =&\frac{1}{(2\pi\imath)^2}\int_0^\infty \mathrm{d}s\exp\left(\frac{\lambda^2}{\nu}s\right)\int\mathrm{d}z_1\exp\left(\frac{\nu t}{2}z_1^2+(x_1-s)z_1\right)\\ &\times\int\mathrm{d}z_2\,(z_1-z_2)\exp\left(\frac{\nu t}{2}z_2^2+(x_2+s)z_2\right). \end{aligned}\tag{2.1}$$

The $\mathrm{d}z_2$ integration over $\alpha_2+\imath\mathbb{R}$ is equal to

$$\begin{aligned}&\int\mathrm{d}z_2\,(z_1-z_2)\exp\left(\frac{\nu t}{2}z_2^2+(x_2+s)z_2\right)\\ &\quad=\frac{\sqrt{2\pi}}{\sqrt{\nu t}}\exp\left(-\frac{(x_2+s)^2}{2\nu t}\right)\\ &\qquad\times\int\mathrm{d}z_2\left(\left[z_1+\frac{x_2+s}{\nu t}\right]-\left[z_2+\frac{x_2+s}{\nu t}\right]\right)\frac{\exp\left(\frac{\nu t}{2}\left[z_2+\frac{x_2+s}{\nu t}\right]^2\right)}{\sqrt{2\pi/(\nu t)}}\\ &\quad=\imath\frac{\sqrt{2\pi}}{(\nu t)^{3/2}}\exp\left(-\frac{(s+x_2)^2}{2\nu t}\right)(\nu t z_1+(x_2+s)). \end{aligned}\tag{2.2}$$

The $\mathrm{d}z_1$ integration over $\alpha_1+\imath\mathbb{R}$ gives

$$\begin{aligned}&\imath\int\exp\left(\frac{\nu t}{2}z_1^2+(x_1-s)z_1\right)\frac{\sqrt{2\pi}}{(\nu t)^{3/2}}\exp\left(-\frac{(s+x_2)^2}{2\nu t}\right)(\nu t z_1+(x_2+s))\,\mathrm{d}z_1\\ &=\imath\frac{2\pi}{\nu t}\exp\left(-\frac{(x_1-s)^2+(x_2+s)^2}{2\nu t}\right)\\ &\quad\times\int\mathrm{d}z_1\frac{1}{\sqrt{2\pi/(\nu t)}}\exp\left(\frac{\nu t}{2}\left[z_1+\frac{x_1-s}{\nu t}\right]^2\right)\left(\left[z_1+\frac{x_1-s}{\nu t}\right]+\frac{x_2-x_1+2s}{\nu t}\right)\\ &=\imath^2\frac{2\pi}{(\nu t)^2}\exp\left(-\frac{(x_1-s)^2+(x_2+s)^2}{2\nu t}\right)(2s+x_2-x_1). \end{aligned}\tag{2.3}$$

Hence,

$$\begin{aligned}&\mathbb{E}[u(t,x_1)u(t,x_2)]\\ &=\int_0^\infty\frac{(2s+x_2-x_1)}{2\pi(\nu t)^2}\exp\left(-\frac{(x_1-s)^2+(x_2+s)^2}{2\nu t}+\frac{\lambda^2 s}{\nu}\right)\mathrm{d}s\\ &=\frac{1}{\pi\nu t}\exp\left(-\frac{x_1^2+x_2^2}{2\nu t}+\frac{(x_1-x_2+t\lambda^2)^2}{4\nu t}\right)\end{aligned}$$

$$
\begin{aligned}
&\times \int_0^\infty \frac{\left(s-\frac{x_1-x_2}{2}\right)}{\nu t} \exp\left(-\frac{1}{\nu t}\left(s-\frac{x_1-x_2+t\lambda^2}{2}\right)^2\right) \mathrm{d}s \\
&= \frac{1}{\pi\nu t}\exp\left(-\frac{x_1^2+x_2^2}{2\nu t}+\frac{(x_1-x_2+t\lambda^2)^2}{4\nu t}\right)\int_{\frac{x_2-x_1-t\lambda^2}{2\sqrt{\nu t}}}^\infty \left(z+\frac{\lambda^2\sqrt{t}}{\sqrt{\nu}}\right)e^{-z^2}\mathrm{d}z \\
&= \frac{1}{\pi\nu t}\exp\left(-\frac{x_1^2+x_2^2}{2\nu t}+\frac{(x_1-x_2+t\lambda^2)^2}{4\nu t}\right) \\
&\quad \times\left(\frac{1}{2}\exp\left(-\frac{(x_1-x_2+t\lambda^2)^2}{4\nu t}\right)+\frac{\lambda^2\sqrt{\pi t}}{2\sqrt{\nu}}\operatorname{erfc}\left(\frac{x_2-x_1-t\lambda^2}{2\sqrt{\nu t}}\right)\right) \\
&= G_\nu(t,x_1)G_\nu(t,x_2) \\
&\quad +\frac{\lambda^2}{4\nu}G_{\nu/2}\left(t,\frac{x_1+x_2}{2}\right)\exp\left(\frac{t\lambda^4-2\lambda^2(x_2-x_1)}{4\nu}\right)\operatorname{erfc}\left(\frac{(x_2-x_1)-t\lambda^2}{2\sqrt{\nu t}}\right). \qquad (2.4)
\end{aligned}
$$

This proves Theorem 1.1. □

3 Third moment: Proof of Theorem 1.2

We first prove two lemmas. For all $\beta\in\mathbb{R}$, $t>0$ and $n\geq 0$, denote

$$\Lambda_n(\beta,t):=\int_0^\infty \exp\left(-\frac{s^2}{t}+\beta s\right)s^n\mathrm{d}s. \qquad (3.1)$$

Lemma 3.1. *For all $t>0$ and $\beta\in\mathbb{R}$,*

$$\Lambda_0(\beta,t)=\frac{\sqrt{\pi t}}{2}\exp\left(\frac{\beta^2 t}{4}\right)\left(2-\operatorname{erfc}\left(\frac{\beta\sqrt{t}}{2}\right)\right), \qquad (3.2)$$

$$\Lambda_1(\beta,t)=\frac{t}{2}+\frac{\sqrt{\pi}t^{3/2}\beta}{4}\exp\left(\frac{\beta^2 t}{4}\right)\left(2-\operatorname{erfc}\left(\frac{\beta\sqrt{t}}{2}\right)\right), \qquad (3.3)$$

and for all $n\geq 2$,

$$\Lambda_n(\beta,t)=\frac{\beta t}{2}\Lambda_{n-1}(\beta,t)+\frac{t(n-1)}{2}\Lambda_{n-2}(\beta,t). \qquad (3.4)$$

In particular, we have that

$$\Lambda_2(\beta,t)=\frac{\beta t^2}{4}+\frac{\sqrt{\pi}t^{3/2}\left(\beta^2 t+2\right)}{8}\exp\left(\frac{\beta^2 t}{4}\right)\left(2-\operatorname{erfc}\left(\frac{\beta\sqrt{t}}{2}\right)\right), \qquad (3.5)$$

$$\Lambda_3(\beta,t)=\frac{t^2}{2}+\frac{\beta^2 t^3}{8}+\frac{\sqrt{\pi}\beta t^{5/2}\left(\beta^2 t+6\right)}{16}\exp\left(\frac{\beta^2 t}{4}\right)\left(2-\operatorname{erfc}\left(\frac{\beta\sqrt{t}}{2}\right)\right). \qquad (3.6)$$

Proof. We first calculate $\Lambda_0(\beta,t)$:

$$
\begin{aligned}
\Lambda_0(\beta,t)&=\int_0^\infty \exp\left(-\frac{1}{t}\left(s-\frac{t\beta}{2}\right)^2+\frac{t\beta^2}{4}\right)\mathrm{d}s \\
&=\sqrt{t}\exp\left(\frac{t\beta^2}{4}\right)\int_0^\infty \exp\left(-\left(z-\frac{\beta\sqrt{t}}{2}\right)^2\right)\mathrm{d}z
\end{aligned}
$$

$$= \sqrt{t}\, \exp\left(\frac{t\beta^2}{4}\right) \frac{\sqrt{\pi}}{2} \left(1 + \operatorname{erf}\left(\frac{\beta\sqrt{t}}{2}\right)\right).$$

As for $\Lambda_1(\beta, t)$,

$$\begin{aligned}
\Lambda_1(\beta, t) &= -\frac{t}{2} \int_0^\infty \left[\left(-\frac{s^2}{t} + \beta s\right)' - \beta\right] \exp\left(-\frac{s^2}{t} + \beta s\right) \mathrm{d}s \\
&= -\frac{t}{2} \int_0^\infty \mathrm{d} \exp\left(-\frac{s^2}{t} + \beta s\right) + \frac{\beta t}{2} \Lambda_0(\beta, t) \\
&= \frac{t}{2} + \frac{\beta t}{2} \Lambda_0(\beta, t),
\end{aligned}$$

which is equal to the right-hand side of (3.3) after simplification. For any $n \geq 2$,

$$\begin{aligned}
\Lambda_n(\beta, t) &= -\frac{t}{2} \int_0^\infty \left[\left(-\frac{s^2}{t} + \beta s\right)' - \beta\right] s^{n-1} \exp\left(-\frac{s^2}{t} + \beta s\right) \mathrm{d}s \\
&= -\frac{t}{2} \int_0^\infty s^{n-1} \mathrm{d} \exp\left(-\frac{s^2}{t} + \beta s\right) + \frac{\beta t}{2} \Lambda_{n-1}(\beta, t) \\
&= -\frac{t}{2} \left(0 - (n-1) \int_0^\infty s^{n-2} \exp\left(-\frac{s^2}{t} + \beta s\right) \mathrm{d}s\right) + \frac{\beta t}{2} \Lambda_{n-1}(\beta, t) \\
&= \frac{t(n-1)}{2} \Lambda_{n-2}(\beta, t) + \frac{\beta t}{2} \Lambda_{n-1}(\beta, t),
\end{aligned}$$

which proves (3.4). In particular,

$$\begin{aligned}
\Lambda_2(\beta, t) &= \frac{\beta t^2}{4} + \left(\frac{t}{2} + \frac{\beta^2 t^2}{4}\right) \Lambda_0(\beta, t), \\
\Lambda_3(\beta, t) &= \frac{t^2}{2} + \frac{\beta^2 t^3}{8} + \left(\frac{3\beta t^2}{4} + \frac{\beta^3 t^3}{8}\right) \Lambda_0(\beta, t).
\end{aligned}$$

Then by (3.2), one obtains (3.5) and (3.6). This completes the proof of Lemma 3.1. □

The next lemma is a corollary of Lemma 3.1. We nevertheless state it separately for the convenience of our application.

Lemma 3.2. *For all $a, b, d \in \mathbb{R}$ and $c > 0$, it holds that*

$$\int_0^\infty (as + b) \exp\left(-cs(s - d)\right) \mathrm{d}s = \frac{\sqrt{\pi}(ad + 2b)}{2\sqrt{c}} \exp\left(\frac{cd^2}{4}\right) \Phi\left(\frac{\sqrt{2c}\, d}{2}\right) + \frac{a}{2c}. \tag{3.7}$$

Now we are ready to prove Theorem 1.2.

Proof of Theorem 1.2. **(1)** Fix $x_1, x_2, x_3 \in \mathbb{R}$ with $x_1 \leq x_2 \leq x_3$, $t > 0$, and $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ with $\alpha_1 > \alpha_2 + \frac{\lambda^2}{\nu} > \alpha_3 + 2\frac{\lambda^2}{\nu}$. Similar to the proof of Theorem 1.1, from (1.8), we have that

$$\begin{aligned}
&\mathbb{E}\left[u(t, x_1) u(t, x_2) u(t, x_3)\right] \\
&\quad = \frac{1}{(2\pi\imath)^3} \int_0^\infty \mathrm{d}s_1 \exp\left(\frac{\lambda^2}{\nu} s_1\right) \int_0^\infty \mathrm{d}s_2 \exp\left(\frac{\lambda^2}{\nu} s_2\right) \int_0^\infty \mathrm{d}s_3 \exp\left(\frac{\lambda^2}{\nu} s_3\right)
\end{aligned}$$

$$
\begin{aligned}
&\times \int \mathrm{d}z_1 \exp\left(\frac{\nu t}{2} z_1^2 + (x_1 - s_1 - s_2) z_1\right) \\
&\times \int \mathrm{d}z_2 \exp\left(\frac{\nu t}{2} z_2^2 + (x_2 + s_1 - s_3) z_2\right) \\
&\times \int \mathrm{d}z_3 (z_1 - z_2)(z_1 - z_3)(z_2 - z_3) \exp\left(\frac{\nu t}{2} z_3^2 + (x_3 + s_2 + s_3) z_3\right). \qquad (3.8)
\end{aligned}
$$

Now we will calculate the triple integral over $\mathrm{d}z_3\mathrm{d}z_2\mathrm{d}z_1$. Recall that $x_1 \le x_2 \le x_3$. The $\mathrm{d}z_3$-integral is equal to

$$
\begin{aligned}
&\int \mathrm{d}z_3 (z_1 - z_2)(z_1 - z_3)(z_2 - z_3) \exp\left(\frac{\nu t}{2} z_3^2 + (x_3 + s_2 + s_3) z_3\right) \\
&\quad = (z_1 - z_2) \int \mathrm{d}z_3 (z_1 - z_3)(z_2 - z_3) \exp\left(\frac{\nu t}{2} z_3^2 + (x_3 + s_2 + s_3) z_3\right) \\
&\quad = (z_1 - z_2) \exp\left(-\frac{(x_3 + s_2 + s_3)^2}{2\nu t}\right) \\
&\qquad \times \int \mathrm{d}z_3 \, \exp\left(\frac{\nu t}{2}\left(z_3 + \frac{x_3 + s_2 + s_3}{\nu t}\right)^2\right) \\
&\qquad\quad \times \left[z_3 + \frac{x_3 + s_2 + s_3}{\nu t} - \left(\frac{x_3 + s_2 + s_3}{\nu t} + z_1\right)\right] \\
&\qquad\quad \times \left[z_3 + \frac{x_3 + s_2 + s_3}{\nu t} - \left(\frac{x_3 + s_2 + s_3}{\nu t} + z_2\right)\right] \\
&\quad = \imath (\nu t)^{-5/2} \sqrt{2\pi} \exp\left(-\frac{(x_3 + s_2 + s_3)^2}{2\nu t}\right) \\
&\qquad \times (z_1 - z_2)\left(-\nu t + (\nu t z_1 + x_3 + s_2 + s_3)(\nu t z_2 + x_3 + s_2 + s_3)\right) \\
&\quad =: F_2(z_2). \qquad (3.9)
\end{aligned}
$$

The $\mathrm{d}z_2$-integral can be obtained in the same way as above

$$
\begin{aligned}
&\int \mathrm{d}z_2 \exp\left(\frac{\nu t}{2} z_2^2 + (x_2 + s_1 - s_3) z_2\right) F_2(z_2) \\
&\quad = \imath (\nu t)^{-5/2} \sqrt{2\pi} \, \exp\left(-\frac{(x_3 + s_2 + s_3)^2}{2\nu t} - \frac{(x_2 + s_1 - s_3)^2}{2\nu t}\right) \\
&\qquad \times \int \mathrm{d}z_2 \exp\left(\frac{\nu t}{2}\left(z_2 + \frac{x_2 + s_1 - s_3}{\nu t}\right)^2\right) \\
&\qquad\quad \times \left[z_1 + \frac{x_2 + s_1 - s_3}{\nu t} - \left(z_2 + \frac{x_2 + s_1 - s_3}{\nu t}\right)\right] \\
&\qquad\quad \times \left[-\nu t + (\nu t z_1 + x_3 + s_2 + s_3)\left(\nu t \left[z_2 + \frac{x_2 + s_1 - s_3}{\nu t}\right] - s_1 + s_2 + 2s_3 - x_2 + x_3\right)\right] \\
&\quad = \imath^2 (\nu t)^{-4} 2\pi (-s_1 + s_2 + 2s_3 - x_2 + x_3) \exp\left(-\frac{(x_3 + s_2 + s_3)^2}{2\nu t} - \frac{(x_2 + s_1 - s_3)^2}{2\nu t}\right) \\
&\qquad \times \left[\nu t + (\nu t z_1 + x_3 + s_2 + s_3)\,(\nu t z_1 + x_2 + s_1 - s_3)\right] \\
&\quad =: F_1(z_1). \qquad (3.10)
\end{aligned}
$$

Similarly, one can calculate the $\mathrm{d}z_1$-integral, which is equal to

$$\begin{aligned}
&\int \mathrm{d}z_1 \exp\left(\frac{\nu t}{2} z_1^2 + (x_1 - s_1 - s_2) z_1\right) F_1(z_1) \\
&\quad = \imath^2 (\nu t)^{-4} 2\pi (-s_1 + s_2 + 2s_3 - x_2 + x_3) \\
&\qquad \times \exp\left(-\frac{(x_3 + s_2 + s_3)^2}{2\nu t} - \frac{(x_2 + s_1 - s_3)^2}{2\nu t} - \frac{(x_1 - s_1 - s_2)^2}{2\nu t}\right) \\
&\qquad \times \int \mathrm{d}z_1 \exp\left(\frac{\nu t}{2}\left(z_1^2 + \frac{x_1 - s_1 - s_2}{\nu t}\right)^2\right) \\
&\qquad\quad \times \left\{\nu t + \left[\nu t\left(z_1 + \frac{x_1 - s_1 - s_2}{\nu t}\right) + s_1 + 2s_2 + s_3 - x_1 + x_3\right]\right. \\
&\qquad\qquad \left. \times \left[\nu t\left(z_1 + \frac{x_1 - s_1 - s_2}{\nu t}\right) + 2s_1 + s_2 - s_3 - x_1 + x_2\right]\right\} \\
&\quad = \imath^3 (\nu t)^{-9/2} (2\pi)^{3/2} \\
&\qquad \times (-s_1 + s_2 + 2s_3 - x_2 + x_3)\,(s_1 + 2s_2 + s_3 - x_1 + x_3)\,(2s_1 + s_2 - s_3 - x_1 + x_2) \\
&\qquad \times \exp\left(-\frac{(x_3 + s_2 + s_3)^2}{2\nu t} - \frac{(x_2 + s_1 - s_3)^2}{2\nu t} - \frac{(x_1 - s_1 - s_2)^2}{2\nu t}\right) \\
&\quad =: F_0(s_1, s_2, s_3).
\end{aligned} \tag{3.11}$$

Hence,

$$\begin{aligned}
&\mathbb{E}\left[u(t, x_1) u(t, x_2) u(t, x_3)\right] \\
&= (\nu t)^{-9/2} (2\pi)^{-3/2} \int_0^\infty \mathrm{d}s_1 \, \exp\left(\frac{\lambda^2 s_1}{\nu}\right) \int_0^\infty \mathrm{d}s_2 \, \exp\left(\frac{\lambda^2 s_2}{\nu}\right) \int_0^\infty \mathrm{d}s_3 \, \exp\left(\frac{\lambda^2 s_3}{\nu}\right) \\
&\quad \times (-s_1 + s_2 + 2s_3 - x_2 + x_3)\,(s_1 + 2s_2 + s_3 - x_1 + x_3)\,(2s_1 + s_2 - s_3 - x_1 + x_2) \\
&\quad \times \exp\left(-\frac{(x_3 + s_2 + s_3)^2}{2\nu t} - \frac{(x_2 + s_1 - s_3)^2}{2\nu t} - \frac{(x_1 - s_1 - s_2)^2}{2\nu t}\right).
\end{aligned} \tag{3.12}$$

Finally, for general $x_i \in \mathbb{R}$, $i = 1, 2, 3$, without ordering, one obtains (1.9) by symmetry.

(2) In this part, we assume $x = x_1 = x_2 = x_3$. Then F_0 defined in (3.11) is equal to

$$\begin{aligned}
F_0(s_1, s_2, s_3) =& \imath^3 (\nu t)^{-9/2} (2\pi)^{3/2} \\
&\times (-s_1 + s_2 + 2s_3)\,(s_1 + 2s_2 + s_3)\,(2s_1 + s_2 - s_3) \\
&\times \exp\left(-\frac{s_1^2 + s_2^2 + s_3^2 + s_1 s_2 - s_1 s_3 + s_2 s_3}{\nu t} - \frac{3x^2}{2\nu t}\right).
\end{aligned} \tag{3.13}$$

Now we are going to apply Lemma 3.1 to evaluate the $\mathrm{d}s_3$-integral. First notice that

$$\begin{aligned}
&(-s_1 + s_2 + 2s_3)\,(s_1 + 2s_2 + s_3)\,(2s_1 + s_2 - s_3) \\
&\quad = (-s_1 + s_2)\,(2s_1 + s_2)\,(s_1 + 2s_2) + 3\left(s_1^2 + 4s_2 s_1 + s_2^2\right) s_3 + 3(s_1 - s_2) s_3^2 - 2s_3^3 \\
&\quad =: a_0 + a_1 s_3 + a_2 s_3^2 + a_3 s_3^3.
\end{aligned} \tag{3.14}$$

Hence,

$$\begin{aligned}\int_0^\infty \mathrm{d}s_3\ \exp\left(\frac{\lambda^2}{\nu}s_3\right) F_0(s_1,s_2,s_3) =& \imath^3(\nu t)^{-9/2}(2\pi)^{3/2}\exp\left(-\frac{3x^2}{2\nu t}-\frac{s_1^2+s_2^2+s_1s_2}{\nu t}\right)\\ &\times\sum_{k=0}^3 a_k\int_0^\infty s_3^k\ \exp\left(-\frac{s_3^2}{\nu t}+\left(\frac{\lambda^2}{\nu}+\frac{s_1-s_2}{\nu t}\right)s_3\right)\mathrm{d}s_3\\ =&\imath^3(\nu t)^{-9/2}(2\pi)^{3/2}\exp\left(-\frac{3x^2}{2\nu t}-\frac{s_1^2+s_2^2+s_1s_2}{\nu t}\right)\\ &\times\sum_{k=0}^3 a_k\,\Lambda_k\left(\frac{\lambda^2}{\nu}+\frac{s_1-s_2}{\nu t},\nu t\right), \end{aligned}\tag{3.15}$$

where the functions $\Lambda_k(\cdot,\cdot)$ are defined in (3.1). After some (tedious) expansion and simplification, we see that

$$\begin{aligned}&\sum_{k=0}^3 a_k\Lambda_k\left(\frac{\lambda^2}{\nu}+\frac{s_1-s_2}{\nu t},\nu t\right)\\ &\quad= \quad b_0+b_1s_2+b_2s_2^2\\ &\qquad-\frac{\sqrt{\pi\nu}\lambda^2t^{3/2}}{4}\exp\left(\frac{(s_1-s_2+\lambda^2t)^2}{4\nu t}\right)\left(t\left(6\nu+\lambda^4t\right)-9\left(s_1+s_2\right)^2\right)\\ &\qquad+\frac{\sqrt{\pi\nu}\lambda^2t^{3/2}}{8}\left(t\left(6\nu+\lambda^4t\right)-9\left(s_1+s_2\right)^2\right)\exp\left(\frac{(s_1-s_2+\lambda^2t)^2}{4\nu t}\right)\operatorname{erfc}\left(\frac{s_1-s_2+\lambda^2t}{2\sqrt{\nu t}}\right)\\ &=:A_1(s_2)+A_2(s_2)+A_3(s_2),\end{aligned}\tag{3.16}$$

where

$$b_0=\nu t\left(2s_1^2+\frac{\lambda^2t}{4}s_1-\nu t-\frac{\lambda^4t^2}{4}\right),\qquad b_1=\nu t\left(5s_1-\frac{\lambda^2t}{4}\right)\quad\text{and}\quad b_2=2\nu t.\tag{3.17}$$

Therefore, the $\mathrm{d}s_2$-integral is equal to

$$\begin{aligned}\int_0^\infty \mathrm{d}s_2\ \exp\left(\frac{\lambda^2}{\nu}s_2\right)\int_0^\infty \mathrm{d}s_3\ \exp\left(\frac{\lambda^2}{\nu}s_3\right)F_0(s_1,s_2,s_3) &= \imath^3(\nu t)^{-9/2}(2\pi)^{3/2}\\ \times\exp\left(-\frac{3x^2}{2\nu t}-\frac{s_1^2}{\nu t}\right)\sum_{k=0}^3\int_0^\infty \mathrm{d}s_2\ \exp\left(-\frac{s_2^2}{\nu t}+\left(\frac{\lambda^2}{\nu}-\frac{s_1}{\nu t}\right)s_2\right)A_k(s_2).&\end{aligned}\tag{3.18}$$

By Lemma 3.1,

$$\begin{aligned}&\int_0^\infty \mathrm{d}s_2\ \exp\left(-\frac{s_2^2}{\nu t}+\left(\frac{\lambda^2}{\nu}-\frac{s_1}{\nu t}\right)s_2\right)A_1(s_2)\\ &\quad=\sum_{k=0}^2 b_k\Lambda_k\left(\frac{\lambda^2}{\nu}-\frac{s_1}{\nu t},\nu t\right)\\ &\quad=\quad\frac{1}{8}\nu^2t^2\left(16s_1+3\lambda^2t\right)\end{aligned}$$

$$+\frac{1}{16}\sqrt{\pi}\lambda^2\nu^{3/2}t^{5/2}\left(15s_1+\lambda^2t\right)\exp\left(\frac{\left(s_1-\lambda^2t\right)^2}{4\nu t}\right)\left(2-\operatorname{erfc}\left(\frac{\lambda^2t-s_1}{2\sqrt{\nu t}}\right)\right). \quad (3.19)$$

Similarly, for the A_2 term, we have that

$$\begin{aligned}
&\int_0^\infty \mathrm{d}s_2\ \exp\left(-\frac{s_2^2}{\nu t}+\left(\frac{\lambda^2}{\nu}-\frac{s_1}{\nu t}\right)s_2\right)A_2(s_2)\\
&\quad=\frac{1}{4}\sqrt{\pi\nu}\lambda^2t^{3/2}\exp\left(\frac{\left(s_1+\lambda^2t\right)^2}{4\nu t}\right)\\
&\qquad\times\int_0^\infty \mathrm{d}s_2\ \exp\left(-\frac{3s_2^2}{4\nu t}+\left(\frac{\lambda^2}{2\nu}-\frac{3s_1}{2\nu t}\right)s_2\right)\left(9\left(s_1+s_2\right)^2-t\left(6\nu+\lambda^4t\right)\right)\\
&\quad=\frac{1}{2}\sqrt{\pi}\lambda^2\nu^{3/2}t^{5/2}\left(3s_1+\lambda^2t\right)\exp\left(\frac{\left(s_1+\lambda^2t\right)^2}{4\nu t}\right), \qquad (3.20)
\end{aligned}$$

where we have applied Lemma 3.1 in the last step. The integration with respect to A_3 is much more complicated. Instead, we claim that for $r>0$,

$$\begin{aligned}
I(r):=&\int_0^r \mathrm{d}s_2\ \exp\left(-\frac{s_2^2}{\nu t}+\left(\frac{\lambda^2}{\nu}-\frac{s_1}{\nu t}\right)s_2\right)A_3(s_2)\\
=&\frac{1}{8}\sqrt{\pi\nu}\lambda^2t^{3/2}\int_0^r \mathrm{d}s_2\ \exp\left(-\frac{s_2^2}{\nu t}+\left(\frac{\lambda^2}{\nu}-\frac{s_1}{\nu t}\right)s_2\right)\left(t\left(6\nu+\lambda^4t\right)-9\left(s_1+s_2\right)^2\right)\\
&\times\exp\left(\frac{\left(s_1-s_2+\lambda^2t\right)^2}{4\nu t}\right)\operatorname{erfc}\left(\frac{s_1-s_2+\lambda^2t}{2\sqrt{\nu t}}\right)\\
=&\frac{1}{16}\lambda^2\nu t^2\Bigg(\sqrt{\pi\nu t}\Bigg[\lambda^2t\Bigg\{-4\exp\left(\frac{\left(s_1+\lambda^2t\right)^2}{4\nu t}\right)\operatorname{erfc}\left(\frac{s_1+\lambda^2t}{2\sqrt{\nu t}}\right)\\
&+5\exp\left(\frac{\left(s_1-\lambda^2t\right)^2}{4\nu t}\right)\left(\operatorname{erfc}\left(\frac{2r+s_1-\lambda^2t}{2\sqrt{\nu t}}\right)+\operatorname{erfc}\left(\frac{\lambda^2t-s_1}{2\sqrt{\nu t}}\right)\right)\Bigg\}\\
&+4\left(3r+\lambda^2t\right)\exp\left(-\frac{3r^2}{4\nu t}+\frac{r\left(2\lambda^2t-6s_1\right)}{4\nu t}+\frac{2\lambda^2s_1t+s_1^2+\lambda^4t^2}{4\nu t}\right)\operatorname{erfc}\left(\frac{-r+s_1+\lambda^2t}{2\sqrt{\nu t}}\right)\Bigg]\\
&+3\sqrt{\pi\nu t}s_1\Bigg[-4\exp\left(\frac{\left(s_1+\lambda^2t\right)^2}{4\nu t}\right)\operatorname{erfc}\left(\frac{s_1+\lambda^2t}{2\sqrt{\nu t}}\right)\\
&+\exp\left(\frac{\left(s_1-\lambda^2t\right)^2}{4\nu t}\right)\operatorname{erfc}\left(\frac{\lambda^2t-s_1}{2\sqrt{\nu t}}\right)\\
&+\exp\left(\frac{\left(s_1-\lambda^2t\right)^2}{4\nu t}\right)\operatorname{erfc}\left(\frac{2r+s_1-\lambda^2t}{2\sqrt{\nu t}}\right)\\
&+4\exp\left(-\frac{3r^2}{4\nu t}+\frac{r\left(2\lambda^2t-6s_1\right)}{4\nu t}+\frac{2\lambda^2s_1t+s_1^2+\lambda^4t^2}{4\nu t}\right)\operatorname{erfc}\left(\frac{-r+s_1+\lambda^2t}{2\sqrt{\nu t}}\right)\\
&-2\exp\left(\frac{\left(s_1-\lambda^2t\right)^2}{4\nu t}\right)\Bigg]\\
&-2t\left[5\sqrt{\pi\nu t}\,\lambda^2\exp\left(\frac{\left(s_1-\lambda^2t\right)^2}{4\nu t}\right)+3\nu-3\nu\exp\left(-\frac{r^2}{\nu t}-\frac{r\left(s_1-\lambda^2t\right)}{\nu t}\right)\right]\Bigg), \qquad (3.21)
\end{aligned}$$

which can be verified directly by differentiating both sides. Because $\mathrm{erfc}(x) \sim x^{-1}\pi^{-1/2}e^{-x^2}$ as $x \to \infty$ and $\lim_{x\to-\infty}\mathrm{erfc}(x) = 2$, we see that

$$\begin{aligned}\lim_{r\to\infty} I(r) =& \frac{1}{16}\lambda^2\nu t^2\Bigg(\sqrt{\pi\nu t}\left[\lambda^2 t\left\{-4\exp\left(\frac{(s_1+\lambda^2 t)^2}{4\nu t}\right)\mathrm{erfc}\left(\frac{s_1+\lambda^2 t}{2\sqrt{\nu t}}\right)\right.\right.\\ &\left.+5\ \exp\left(\frac{(s_1-\lambda^2 t)^2}{4\nu t}\right)\mathrm{erfc}\left(\frac{\lambda^2 t-s_1}{2\sqrt{\nu t}}\right)\right\}+0\Bigg]\\ &+3\sqrt{\pi\nu t}s_1\Bigg[-4\exp\left(\frac{(s_1+\lambda^2 t)^2}{4\nu t}\right)\mathrm{erfc}\left(\frac{s_1+\lambda^2 t}{2\sqrt{\nu t}}\right)\\ &+\exp\left(\frac{(s_1-\lambda^2 t)^2}{4\nu t}\right)\mathrm{erfc}\left(\frac{\lambda^2 t-s_1}{2\sqrt{\nu t}}\right)\\ &+0+0-2\exp\left(\frac{(s_1-\lambda^2 t)^2}{4\nu t}\right)\Bigg]\\ &-2t\left[5\sqrt{\pi\nu t}\,\lambda^2\exp\left(\frac{(s_1-\lambda^2 t)^2}{4\nu t}\right)+3\nu-0\right]\Bigg).\end{aligned} \tag{3.22}$$

Therefore,

$$\begin{aligned}&\int_0^\infty \mathrm{d}s_2\ \exp\left(-\frac{s_2^2}{\nu t}+\left(\frac{\lambda^2}{\nu}-\frac{s_1}{\nu t}\right)s_2\right)A_3(s_2)\\ &=\frac{1}{16}\lambda^2\nu t^2\Bigg(\sqrt{\pi\nu}\,\lambda^2 t^{3/2}\left\{-4\exp\left(\frac{(s_1+\lambda^2 t)^2}{4\nu t}\right)\mathrm{erfc}\left(\frac{s_1+\lambda^2 t}{2\sqrt{\nu t}}\right)\right.\\ &\left.+5\ \exp\left(\frac{(s_1-\lambda^2 t)^2}{4\nu t}\right)\mathrm{erfc}\left(\frac{\lambda^2 t-s_1}{2\sqrt{\nu t}}\right)\right\}\\ &+3\sqrt{\pi\nu t}s_1\Bigg[-4\ \exp\left(\frac{(s_1+\lambda^2 t)^2}{4\nu t}\right)\mathrm{erfc}\left(\frac{s_1+\lambda^2 t}{2\sqrt{\nu t}}\right)\\ &+\exp\left(\frac{(s_1-\lambda^2 t)^2}{4\nu t}\right)\mathrm{erfc}\left(\frac{\lambda^2 t-s_1}{2\sqrt{\nu t}}\right)-2\exp\left(\frac{(s_1-\lambda^2 t)^2}{4\nu t}\right)\Bigg]\\ &-2t\left[5\sqrt{\pi\nu t}\,\lambda^2\exp\left(\frac{(s_1-\lambda^2 t)^2}{4\nu t}\right)+3\nu\right]\Bigg).\end{aligned} \tag{3.23}$$

Finally, combining all these three integrals, we see that

$$\begin{aligned}&\exp\left(\frac{\lambda^2}{\nu}s_1\right)\int_0^\infty \mathrm{d}s_2\ \exp\left(\frac{\lambda^2}{\nu}s_2\right)\int_0^\infty \mathrm{d}s_3\ \exp\left(\frac{\lambda^2}{\nu}s_3\right)F_0(s_1,s_2,s_3)\\ &=\imath^3\exp\left(-\frac{3x^2}{2\nu t}\right)\\ &\quad\times\left[\frac{\pi^2\lambda^2}{\sqrt{2}\,\nu^3 t^2}\left(3s_1+\lambda^2 t\right)\exp\left(\frac{\lambda^4 t}{\nu}-\frac{3\left(s_1-\lambda^2 t\right)^2}{4\nu t}\right)\left(2-\mathrm{erfc}\left(\frac{s_1+\lambda^2 t}{2\sqrt{\nu t}}\right)\right)\right.\end{aligned}$$

$$+\frac{\pi^2\lambda^2\left(3s_1-\lambda^2 t\right)}{\sqrt{2}\,\nu^3 t^2}\exp\left(\frac{t\lambda^4}{3\nu}-\frac{3\left(s_1-\lambda^2 t/3\right)^2}{4\nu t}\right)\left(2-\operatorname{erfc}\left(\frac{\lambda^2 t-s_1}{2\sqrt{\nu t}}\right)\right)$$
$$+2^{5/2}(\nu t)^{-5/2}\pi^{3/2}s_1\exp\left(\frac{\lambda^4 t}{4\nu}-\frac{\left(s_1-\lambda^2 t/2\right)^2}{\nu t}\right)\Bigg]$$
$$=:\Theta(s_1). \tag{3.24}$$

Therefore, the third moment is equal to

$$\mathbb{E}\left[u(t,x)^3\right]=\frac{1}{(2\pi\imath)^3}\int_0^\infty \Theta(s_1)\mathrm{d}s_1. \tag{3.25}$$

Because $2-\operatorname{erfc}(x)=2\Phi\left(\sqrt{2}x\right)$ and thanks to Lemma 3.2,

$$\begin{aligned}
&\frac{1}{(2\pi)^3}\exp\left(-\frac{3x^2}{2\nu t}\right)\int_0^\infty 2^{5/2}(\nu t)^{-5/2}\pi^{3/2}s_1\exp\left(\frac{\lambda^4 t}{4\nu}-\frac{\left(s_1-\lambda^2 t/2\right)^2}{\nu t}\right)\mathrm{d}s_1\\
=&\frac{1}{\sqrt{2}\pi^{3/2}(\nu t)^{5/2}}\exp\left(-\frac{3x^2}{2\nu t}\right)\int_0^\infty s_1\exp\left(-\frac{s_1\left(s_1-\lambda^2 t\right)}{\nu t}\right)\mathrm{d}s_1\\
=&\frac{1}{(2\pi\nu t)^{3/2}}\exp\left(-\frac{3x^2}{2\nu t}\right)+\frac{\lambda^2}{2\sqrt{2}\pi\nu^2 t}\exp\left(\frac{\lambda^4 t}{4\nu}-\frac{3x^2}{2\nu t}\right)\left(1-\Phi\left(-\lambda^2\sqrt{\frac{t}{2\nu}}\right)\right),
\end{aligned} \tag{3.26}$$

one can obtain (1.10) after some simplification.

(3) By the mean-value theorem, there exit two constants a, b in the range given by (1.11) such that

$$\begin{aligned}
\mathbb{E}[u(t,x)^3]=&\frac{b\lambda^2}{2^{5/2}\pi\,\nu^3 t^2}\exp\left(\frac{\lambda^2 t}{4\nu}-\frac{3x^2}{2\nu t}\right)\int_0^\infty\left(3s_1+\lambda^2 t\right)\exp\left(-\frac{3s_1(s_1-2\lambda^2 t)}{4\nu t}\right)\mathrm{d}s_1\\
&+\frac{a\lambda^2}{2^{5/2}\pi\,\nu^3 t^2}\exp\left(\frac{\lambda^2 t}{4\nu}-\frac{3x^2}{2\nu t}\right)\int_0^\infty\left(3s_1-\lambda^2 t\right)\exp\left(-\frac{s_1(3s_1-2\lambda^2 t)}{4\nu t}\right)\mathrm{d}s_1\\
&+\frac{1}{(2\pi\nu t)^{3/2}}\exp\left(-\frac{3x^2}{2\nu t}\right)\\
&+\frac{\lambda^2}{2\sqrt{2}\pi\nu^2 t}\exp\left(\frac{\lambda^4 t}{4\nu}-\frac{3x^2}{2\nu t}\right)\left(1-\Phi\left(-\lambda^2\sqrt{\frac{t}{2\nu}}\right)\right).
\end{aligned} \tag{3.27}$$

Then one can apply Lemma 3.2 to evaluate these two integrals. This completes the whole proof of Theorem 1.2. □

Acknowledgements

L.C. would like to thank *Robert Dalang* and *Davar Khoshnevisan* for many helpful comments. Khoshnevisan asked L.C. in 2014 whether one could obtain an explicit formula for the third moment, which motivated the current study. There was no much progress on this problem until when L.C. was vising the *Simons Center for Geometry and Physics* for the conference – *Stochastic Partial Differential Equations* (May 16-20, 2016), *Ivan Corwin* pointed out to L.C. the transform from (1.6) to (1.8). This became the starting point of the whole calculation in this paper. Here L.C. would like to express his sincere gratitude to him. Finally, L.C. would also like to thank the organizer *Martin Hairer* for the wonderful conference.

References

[1] Bertini, Lorenzo and Nicoletta Cancrini. The stochastic heat equation: Feynman-Kac formula and intermittence. *J. Statist. Phys.*, 78(5-6):1377–1401, 1995.

[2] Borodin, Alexei and Ivan Corwin. Moments and Lyapunov exponents for the parabolic Anderson model. *Ann. Appl. Probab.,* 24 (2014), no. 3, 1172–1198.

[3] Carmona, René A. and Stanislav A. Molchanov. Parabolic Anderson problem and intermittency. *Mem. Amer. Math. Soc.*, 108(518), 1994.

[4] Chen, Le and Robert C. Dalang. Moments and growth indices for nonlinear stochastic heat equation with rough initial conditions. *Ann. Probab.* Vol. 43, No. 6, 3006–3051, 2015.

[5] Chen, Le and Kunwoo Kim. Nonlinear stochastic heat equation driven by spatially colored noise: moments and intermittency. Preprint arXiv:1510.06046, 2015.

[6] Chen, Le and Kunwoo Kim. On comparison principle and strict positivity of solutions to the nonlinear stochastic fractional heat equations. *Ann. Inst. Henri Poincaré Probab. Stat.*. to appear, 2016.

[7] Chen, Le, Yaozhong Hu and David Nualart. Two-point correlation function and Feynman-Kac formula for the stochastic heat equation. *Potential Anal.*, to appear, 2016.

[8] Chen, Xia. Precise intermittency for the parabolic Anderson equation with an $(1 + 1)$-dimensional timespace white noise. *Ann. Inst. Henri Poincaré Probab. Stat.* 2015, Vol. 51, No. 4, 1486–1499.

[9] Conus, Daniel and Davar Khoshnevisan. On the existence and position of the farthest peaks of a family of stochastic heat and wave equations. *Probab. Theory Related Fields*, 152(3-4) (2012) 681–701.

[10] Foondun, Mohammud and Davar Khoshnevisan. Intermittence and nonlinear parabolic stochastic partial differential equations. *Electron. J. Probab.* 14(21) (2009) 548–568.

[11] Huang, Jingyu, Khoa Lê and David Nualart. Large time asymptotics for the parabolic Anderson model driven by spatially correlated noise. Preprint arXiv:1509.00897v3, 2015.

[12] Huang, Jingyu, Khoa Lê and David Nualart. Large time asymptotics for the parabolic Anderson model driven by space and time correlated noise. Preprint arXiv:1607.00682, 2016.

[13] Walsh, John B. *An Introduction to Stochastic Partial Differential Equations.* In: École d'été de probabilités de Saint-Flour, XIV—1984, 265–439. Lecture Notes in Math. 1180, Springer, Berlin, 1986.